

**Central Bureau for Astronomical Telegrams**  
**INTERNATIONAL ASTRONOMICAL UNION**

Mailstop 18, Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.  
 IAUSUBS@CFA.HARVARD.EDU or FAX 617-495-7231 (subscriptions)  
 CBAT@CFA.HARVARD.EDU (science)  
 URL <http://www.cfa.harvard.edu/iau/cbat.html> ISSN 0081-0304  
 Phone 617-495-7440/7244/7444 (for emergency use only)

*COMET C/2008 E3 (GARRADD)*

G. J. Garradd reports his discovery of a comet on images taken with the 0.5-m Uppsala Schmidt telescope at Siding Spring (discovery observation tabulated below), the object being diffuse and 20'' in diameter on images from Mar. 7.8 UT. Following posting on the 'NEOCP' webpage, E. Guido and G. Sostero (Castellammare di Stabia, Italy) write that they obtained CCD images remotely using a 0.25-m f/3.4 reflector near Moorook, Australia, on Mar. 8.7; thirty co-added 60-s exposures show the comet to have cometary appearance, with a coma diameter of nearly 10'' with a strong central condensation and a broad tail 18'' long toward the north.

2008	UT	$\alpha_{2000}$	$\delta_{2000}$	Mag.
Mar.	5.77805	19 <sup>h</sup> 35 <sup>m</sup> 19 <sup>s</sup> .03	-41°44'05''.8	17.8

The available astrometry, the following preliminary parabolic orbital elements, and an ephemeris appear on *MPEC* 2008-E70.

$$\begin{array}{rcl}
 T = 2008 \text{ Oct. } 31.240 \text{ TT} & \omega = 229.143 & \\
 & \Omega = 106.195 & \\
 q = 5.26127 \text{ AU} & i = 106.590 & \left. \vphantom{\begin{array}{l} \omega \\ \Omega \\ i \end{array}} \right\} 2000.0
 \end{array}$$

*NOVA CYGNI 2008*

S. Nakano, Sumoto, Japan, reports the discovery by H. Kaneda (Sapporo, Japan) of an apparent nova (mag  $8.2 \pm 0.3$ ) on unfiltered CCD frames taken on Mar. 7.801 UT using a telephoto lens, giving the position of the variable as  $\alpha = 19^{\text{h}}58^{\text{m}}33^{\text{s}}.4$ ,  $\delta = +29^{\circ}52'04''$  (equinox 2000.0). Nakano forwards the following precise position and figures for the nova from a CCD exposure on Mar. 8.716 by K. Kadota (Ageo, Japan, 0.25-m reflector): 33<sup>s</sup>.39, 06''5, mag 7.4. Numerous additional apparent independent discoveries were reported on *CBET* 1291.

D. Nogami and J. Kuriyama, Kyoto University; and I. Iwata, National Astronomical Observatory of Japan, write that a low-resolution ( $R \sim 500$ ) spectrum (range 400–800 nm), taken of the new variable on Mar. 8.794 UT with a 1.88-m telescope (+ KOOLS) at the Okayama Astrophysical Observatory, shows a blue continuum with strong Balmer and Fe II lines having prominent P-Cyg profiles. The FWHM of the emission component of H $\alpha$  is  $\sim 1000$  km/s, and the minimum of the H $\alpha$  absorption is blueshifted from the emission peak by  $\sim 880$  km/s. These features suggest that this object is indeed a classical nova before or around maximum.